Sets And Functions

Sr	Questions	Answers Choice
1	In cartesian plane, vertically line is called	A. x - axis B. y - axis C. point D. origin
2	The table which gives the frequency of each score is called	A. logarithmic table B. grouped table C. ungrouped table D. frequency table
3	$\bar{x}$ is called	A. arithmetic mean B. mode C. median D. group data
4	A running total of class frequency is called	A. histogram B. cumulative frequency C. data D. call interval
5	According of grouped data formula of arithmetic mean	A. $x = \sum fx/\sum f$ B. $x = \sum fx/\sum n$ C. $x = \sum fx/\sum x$ D. $x = \sum x/\sum n$
6	The number of time each value appears in the date is called	A. polygon B. frequency C. histogram D. frequency table
7	In cartesian plane point 'O' is called	A. column B. row C. origin D. axis
8	First elements of ordered pairs Is called:	A. column B. row C. point D. origin
9	The formula used to convert Celsius temperature (°C)into Fahrenheit temperature (°F) is"	A. °C = 9/5(°F -32) B. °C = 5/9(°C +32) C. °C = 5/9(°F + 32) D. °C = 5/9(°F -32)
10	1 miles =?	A. 1.5 km, B. 1.6 km, C. 6.1 km, D. 5.1 km,
11	If a group of students get marks from 20% to 10% their frequency will be	A. 10 B. 15 C. 20 D. 5
12	A set of two elements, listed in a specific order is called	A. unordered pairs B. ordered pairs C. cartesian D. rectangular
13	The coordinates of origin are	A. (1,0) B. (0,1) C. (0,0) D. (1,1)
14	In Cartesian plane the horizontal line XOX' IS	A. x - axis B. y - axis C. origin D. co-ordinate axis
15	The formula used to convert Fahrenheit temperature (°F) into celsius temperature (°C) is:	A. °F = (9/5 X ° c) + 33 B. °F = (9/5 X ° c) + 32 C. °F = 9/5(°F - 32) D. °F = 9/5(°F + 32)